



UNIVERSIDAD AUTÓNOMA DEL
ESTADO DE MORELOS




Molecular phylogeny of the species of the genus *Ilyodon* (Goodeidae)

Rosa Gabriela Beltrán López, Ignacio Doadrio
and Omar Domínguez Domínguez

BACKGROUND

- Volcanic and tectonic activities
- Miocene, Pliocene, Pleistocene
- continues cycles of communication and isolation over the past million years
- High richness of freshwater fishes species (near to 100 spp.)
- High number of endemism (near to 70%)

Central Mexico



(Barbour, 1973; Smith, 1980; Ferrari et al. 2000).

FAMILY GOODEIDAE

- 20 genera
- 42 species
- Subfamily Goodeinae





I. whitei (Meek, 1904)



I. furcidens (Jordan y Gilbert, 1882)



I. amecae (Radda, 1989)



I. lennoni (Meyer y Foerster, 1983)



I. xantusi (Hubbs y Turner, 1939)



I. cortesae (Paulo-Maya y Trujillo-Jimenéz, 2000)

Controversies in the number of species that comprise the genus *Ilyodon*

- **Turner y Grosse (1980)** compare 5 allozyme loci and proposed that *I. furcidens* and *I. xantusi*, are not different species, but have different trophic morphotypes
- **Doadrio and Domínguez (2004)** recognized three species: *I. whitei*, *I. furcidens* and *I. amecae* (0.01-1.7%).
- **Webb et al. (2004)** recognized two species: *I. whitei* and *I. furcidens* (0.8%).
- **Miller et al. (2005)** recognized only to *I. whitei* and *I. furcidens*
- **Domínguez-Domínguez et al. (2010)** they described to *I. whitei* and *I. furcidens* as paraphyletic species, because *I. whitei* is grouped like a polytomy unresolved that included to *I. lennoni*, *I. cortesae* and at least one population of *I. furcidens*

- The present study represent the first one in which it's include samples of all distribution range of species of genus *Ilyodon* and the first that include mitochondrial and nuclear loci.

- The aims of this work were:

1. Infer the genus evolutionary history and relationships between all recognized species
2. Establish the time and the way of diversification of main clades recovered


Also, considering that *Ilyodon whitei* is catalogued in the IUCN Red List of Threatened Species (IUCN, 2015)

and another specie is in the NOM-059 (SEMARNAT, 2010)

result important clear up the phylogenetic relationships of the species that composed the genus *Ilyodon*, to establish appropriate conservation programs.

METHODS

84 individuals



DNA Extraction

PCR

Edition & Alignment

Were amplified fragments of three mitochondrial genes:
Cytochrome b (*Cytb*)
Subunit I of Cytochrome Oxidase I (*COI*)
(*Dloop*) of the control region
nuclear gene B-actin (*RAG1*)
4032 pb

Partition homogeneity test in PAUP was conducted to evaluate the significance of conflict among data sets, using 1000 resampling of characters.

Model selection based on Akaike information criterion (AIC) and optimal partition setting analysis were performed using PartitionFinder v1.1.0 (Lanfear et al. 2012). We treated each gene as a single locus.

Haplotype network


Gene trees were constructed using Bayesian inference (BI): Mr. Bayes and Maximum likelihood (RAxML)

Divergence times estimation: (Cytb) molecular clock calibrated with the mutation rate calculated for the teleosts fishes


Uncorrected genetic distances “p” with Cytb and COI (Mega v5.1)

AMOVA: how is partitioned the genetic variance at various hierarchical levels
Cytb
COI

RESULTS




Cytb
One to 12 mutational steps




COI

One to nine mutational steps




Dloop

One to 24 mutational steps





Beta-actin One to 7 mutational steps




RAG1 One to 3 mutational steps

Phylogenetic tree with mtDNA


Bayesian posterior probabilities/Bootstrap probabilities





Phylogenetic tree with mtDNA+nDNA

Bayesian posterior probabilities/Bootstrap probabilities





Divergence time between
Xenotaenia and *Ilyodon*
Miocene epoch

Divergence time between
Populations of Ameca and
Armeria river and the rest of
population

Between Pliocene and Pleistocene
epoch (1.99 Mya)

while the isolation of the
populations of the Coahuayana
river diverged of the populations
of Balsas basin in Pleistocene
epoch (0.98 Mya)





Cytb: 0.9-2.1%
 COI: 0.6-1.2%
 Between clades

	CLADO IA	CLADO IB	CLADO IC	CLADO ID	CLADO IE
<i>Cytb/COI</i>					
CLADO IA	0.00/0.001	0.009	0.008	0.010	0.011
CLADO IB	0.016	0.005/0.005	0.009	0.012	0.012
CLADO IC	0.021	0.019	0.005/0.00	0.006	0.007
CLADO ID	0.014	0.015	0.019	0.003/0.003	0.006
CLADO IE	0.015	0.015	0.017	0.009	0.004/0.001


0.02 subs/site

Analyses of molecular variance with two different grouping (populations of each hydrological basin and populations according with recovered clades) for *cytb* and *COI*.

Groups	df	Sum of squares	Variance components	% Variation
<i>Cytb</i> (grouping with hydrological basin)				
Among groups	4	6.98	0.03 Va	7.11
Among populations within groups	20	19.15	0.20 Vb	41.24
Within populations	70	17.93	0.25 vc	51.65
F_{ST} : 0.48 ($p < 0.001$); F_{CT} : 0.07 (p ns); F_{SC} : 0.44 ($p < 0.001$)				
<i>Cytb</i> (grouping with recovered clades)				
Among groups	4	212.75	3.00 Va	72.09
Among populations within groups	19	48.84	0.62 Vb	15.04
Within populations	66	35.45	0.53 vc	12.88
F_{ST} : 0.87 ($p < 0.001$); F_{CT} : 0.72 ($p < 0.001$); F_{SC} : 0.53 ($p < 0.001$)				
<i>COI</i> (grouping with hydrological basin)				
Among groups	3	9.85	0.15 Va	29.95
Among populations within groups	21	18.70	0.22 Vb	41.85
Within populations	64	9.63	0.15 vc	28.20
F_{ST} : 0.71 ($p < 0.001$); F_{CT} : 0.29 ($p < 0.001$); F_{SC} : 0.59 ($p < 0.001$)				
<i>COI</i> (grouping with recovered clades)				
Among groups	3	63.21	1.63 Va	85.31
Among populations within groups	16	7.00	0.08 Vb	4.52
Within populations	44	8.56	0.19 vc	10.17
F_{ST} : 0.89 ($p < 0.001$); F_{CT} : 0.85 ($p < 0.001$); F_{SC} : 0.30 ($p < 0.001$)				

DISCUSSION

Different species mixed in topologies
(Domínguez-Domínguez et al., 2010)



Genetic distances between two clades of *I. whitei*: 1.6% (*Cytb*), 0.9% (*COI*).


Intraspecific genetic distance
(0.4% *Cytb*, 0.1% *COI*)

I. xantusi, *I. lennoni* and *I. cortesae* are not valid species in this work, are only different trophic morphotypes:

- Mixed in topologies with *I. whitei* and *I. furcidens*
- Small genetic distances
- Shared haplotypes
- Same geographic distribution of *I. furcidens* and *I. whitei*


Recent diversification

- Pliocene and Pleistocene
- Clade IC with the rest coincide with the emergence of the Trans-Mexican-Volcanic Belt



Balsas basin

- High genetic structure
- High, middle and low Balsas basin
- Don't share haplotypes between them
- Independent clades
- Genetic distances (1.5-1.6% Cytb; 0.9-1.2% COI)
- Between Ameca, Armeria and Mascota (0.2-0.7% Cytb; 0% for COI).
- Other species as *Notropis*



Conservation

- Five clades with geographic correspondence
 - Five genetically different groups
 - Different units for conservation
 - *I. whitei* (IUCN, 2015)
 - *I. furcoidens* (NOM-059)



Conclusions

- Five phylogenetic groups have geographic correspondence
- Diversification of genus *Ilyodon* in Pliocene and Pleistocene epoch
- *I. lennoni* and *I. cortesae* should be recognized as a synonym of *I. whitei*
- *I. amecae*, *I. furcidens* and *I. whitei*: Incomplete Lineages Sorting (ILS) or incipient species
- We recommended an extended work especially for populations of Balsas basin.

Acknowledgements

- All people tha help me with the collect of individuals
- Laboratory of Molecular systematic, School of Biology,
Universidad Autónoma del Estado de Morelos
- Laboratory of Molecular systematic Aquatic Biology
Department. Universidad Michoacana de San Nicolás de
Hidalgo. Michoacán. México
- Department of Biodiversity and Evolutionary Biology. National
Museum of Natural Sciences. Madrid. Spain

¡ Thank you very much for your
attention !